antipyrine metabolism.

Erixon-Carlquist E., Hansson T., Lundstrom J.:
Metabolism of raclopride in different species.

Evans G.L., Ayrton J., Grossi P., Wilson K.:
Extra-hepatic metabolism of lacidipine in the rat.

Ferrarese A., Celeste S., Marini P.P., Caccia S.:
Hepatic microsomal metabolism of the potential memory-enhancing agent CL 275, 838 to its desbenzyl derivative.

Ferrari L., Kremers P., Batt A.M., Gielen J.E., Siest G.:
Differential effects of human recombinant interleukin-1B on cytochrome P-450 dependent activities in rats in vitro and in vivo.

Ferrari M.P., Caselli G.F., Clavenna G., Canali S., Giachetti G., Zanolo G.:
Metabolic pattern of [14C]-rispenzepine. HCl in rat, dog and human.

Filaquer C., Peirone M.H., Buchan P., Guyomard C.:
Interspecies metabolic study of adapalene (CD271) in hepatocyte culture.

Fischer V.:
Polymorphic drug metabolism and its consequences for drug development.

Foliot A., Beaune P.:
Effects of microsomal enzyme inducers on glutathione S-transferase isoenzymes in liver of rats and hamsters.

Forster A., Fournel-Gigleux S., Burchell B., Bock K.W.:
Different isozymed of UDP-glucuronosyltransferase (UGT) catalyze high and low affinity paracetamol-UGT activity in rat and man.

Fournel-Gigleux S., Ouzzine M., Pillot T., Pritchard M., Taguchi T., Magdalou J., Siest G., Burchell B.:
Heterologous expression of human UDP-glucuronosyl transferase cDNAs in

2028897196

Source: https://www.industrydocuments.ucsf.edu/docs/hzcl0000

glucuronidation of phenols in rat as well as in man.

Guedes Y., Berthou F., Hourmant N., Le Garrec J., Ratanasavanh D.:
Metabolism study of bromazepam in rat.

Guengerich F.P.:
New vistas on P-450.

Hamdoune M., Mounie J., Goudonnet H., Magdalou J., Escousse A., Truchot R.C.:
Regulation by thyroid status of glucuronidation of flurbiprofen enantiomers.

Hansen K.T., Christensen J.V., Stentoft K., Sorensen L., Kristensen E., Young C.:
Disposition and biotransformation of the novel dopamine D-1 receptor antagonist, NNC 01-0687, in laboratory animals.

Hansen K.T.:
Hepatic uptake and biliary excretion of the neuroprotective drug, NBQX, in the isolated perfused rat liver.

Hayes K., Chenery R., Houston J.B.:
Effect of albumin on phenytoin clearance by hepatocyte suspensions.

Hibert M.F.:
Three dimensional structure of G protein-coupled receptors and drug design.

Humphrey M.J.:
The role of metabolism studies in drug discovery.

Huszka M., Teichmann F.:
Biotransformation of restacorin in the rat and some properties of probable metabolites.

Ioannides C., Cheung Y-L., Wilson J., Lewis D.F.V., Gray I.J.B.:
Mechanisms that may account for the marked difference in carcinogenic potential between 2-acetylaminofluorene and its 4-isomer.

2028897497

Source: https://www.industrydocuments.ucsf.edu/docs/hzcl0000

eukaryotic and prokaryotic cells.

Fracasso C., Anelli M., Ferrarese A., Garattini S., Caccia S.:
Effect of SKF-525A on the brain levels and indole depleting action of D-fenfluramine in rats: hepatic metabolism is not a determinant of its neurochemical action on the serotoninergic system.

Fuming L., Kishore B.K., Tulkens P.M.:
A stable model of anesthetised and infused rats for single dose pharmacokinetic studies.

Funck-Brentano C.:
Active metabolites of cardiovascular drugs.

Galtier P., Kaddouri M., Eeckhoutte C., Sutra J.F., Alvinerie M.:
Hepatic microsomal metabolism of progesterone in sheep.

Gescher A.:
The role of metabolism in the efficacy and toxicity of some novel anticancer drugs.

Gires P., Piguet V., Martin S., Chevalier P., Berteaud-Moantagu M.R., Martinet M., Chapelle P., Frydman A.:
Effect of H2 antagonists on the handling and differential secretion of RP49532 (Girodazole) by isolated perfused rat kidney.

Grabe M., Tegtmeier M., Hahnemann B., Legrum W.:
Specificity of coumarin derivatives towards hepatic and renal coumarin 7-hydroxylase of male DBA/2J-Mice.

Grosa G., Biglino G.:
In vitro metabolism of 7(1,3-thiazolidin-2-ylmethyl) theophylline: evidence for a monooxygenase-dependent step in 1,3-thiazolidine cleavage.

Gschaidmeier H., Beck S., Seidel A., Fournel-Gigleux S., Burchell B., Bock K.W.:
Different isozymes of UDP-glucuronosyltransferase (UGT) are responsible for

2028897198

Source: https://www.industrydocuments.ucsf.edu/docs/hzcl0000

Leemann T., Kondo M., Transon C., Bonnabry P., Dayer P.:
Cytochrome P-450TB (CYP2C subfamily): the common site of NSAIDS oxidation by human liver.

Leemann T., Dayer P.:
Q-DIPS: towards systematic predictions of drug metabolism interactions from in vitro data.

Lock E.A.:
Toxic reactive metabolites in the kidney.

Lopez-Garcia M.P., Valadon P., Dansette P.M., Beaune P., Guengerich F.P., Mansuy D.:
Human cytochromes P-450 2C9 and 2C10 (expressed in yeast) are responsible for metabolic activation of tienilic acid and its isomer.

2028897199

Source: https://www.industrydocuments.ucsf.edu/docs/hzcl0000

Johansson B.:
Human blood mononuclear leucocytes as a model tissue for biotransformation and immuno suppression studies of immunomodulators.

Jugert F.K., Agarwal R., Mukhtar H., Merk H.F.:
Induction of cytochrome P-450 subfamilies 1A, 2B, 2E and 3A in murine skin by dexamethasone.

Jugert F.K., Frankenberg S., Junginger H., Merk H.F.:
Multiple cytochrome P-450 isozymes are present in human keratinocytes.

Kainz A., Gescher A., Chipman J.K.:
Toxicity of 1,1- dichloroethene (vinylidenchloride, DCE) and its metabolites towards mouse hepatocytes in suspensions.

Kehr J.:
Microdialysis in vivo opens up new dimensions in pharmacokinetics.

Kohl C., Schiller C.D., Gescher A., Farmer P.B., Bailey E.:
The hepatocarcinogen acetoxime (ACO) is metabolized to the potent genotoxicant propane-2-nitronate (P2N).

Kopecky J., Vicek J., Kvetina J.:
Brain and red blood cells lithium bioavailability during the experimental chronic renal failure in rats.

Korolenko T.A., Vereschagin E.I., Safina A.F., Urazgaliyev K., Sandula A.F.:
Receptor-mediated uptake of lysosomotropic glycans yeast polysaccharides by liver cells in vivo and in vitro.

Langouet S., Morel F., Fardel O., Meyer D., Ketterer B., Guillouzo A.:
Effect of phenobarbital, 3-methylcholanthrene, 1-2-dithiole 3-thione and oltipraz on the expression and activity of rat glutathione S-transferase subunits in cultured hepatocytes.

2028897200

Source: https://www.industrydocuments.ucsf.edu/docs/hzcl0000